arXiv:1101.0660v1 [quant-ph] 4 Jan 2011

Quantum contextuality as security check in quantum cryptography using intraparticle entanglement

S. Adhikari*
Institute of Physics, Bhubaneswar, India

Dipankar Home†
CAPSS, Dept. of Physics, Bose Institute, Salt Lake, Kolkata-700091, India

A. S. Majumdar‡
S. N. Bose National Centre for Basic Sciences, Salt Lake, Kolkata 700 098, India

A. K. Pan§
Dept. of Physics, Busan, Korea

(Dated: January 5, 2011)

In this work, we present a quantum key distribution scheme which exploits quantum mechanical violation of non-contextuality for checking the security of the protocol. In this scheme the sender encodes the key using path-spin intraparticle entanglement of a single particle. The receiver decodes the information using a combination of beam splitters and phase shifters in conjunction with Stern-Gerlach measuring devices. Distinctive features of this protocol as compared to the BB84 and E91 protocols are discussed.

PACS numbers:

INTRODUCTION

Quantum key distribution (QKD) protocols [1] allow two distant parties, traditionally called Alice and Bob, to produce a shared random bit string consisting of 0's and 1's known only to them, which can be used as a key to encrypt and decrypt messages. Based on the fundamental principles such as no-cloning principle [2], in quantum physics, QKD provides an unconditionally secure way to distribute random keys through insecure channels. In 1984, the well known Bennett-Brassard (BB84) protocol was first proposed [3]. The first entanglement based QKD protocol was proposed by Ekert in 1991 [11], where violation of a local realist inequality is used to check the security of QKD, i.e., if the shared state between Alice and Bob violates Bell's inequality [12], then it is possible to generate a secure key. Subsequently, there have been several important theoretical improvements and experimental demonstrations of BB84 and other QKD protocols were discussed in [4–10]. In this connection, the violation of the Bell's inequality was first monitored in an experiment by Jennewein et. al.[13], but no quantitative measure of security was derived from the observed violation. Later, Ling et. al. [14] have proposed an experiment on entanglement based QKD in which the violation of Bell-CHSH inequality is used to also quantify the degree of security.

In this work, we provide a quantum key distribution scheme in which path-spin entanglement in a single particle (also known as intraparticle entanglement) is exploited. In the first phase (key distribution phase), our protocol proceeds like the BB84 protocol with a notable difference in the preparation of the state. In the BB84 protocol, among the four prepared states, two are classical states (bits), and the remaining two are quantum states (superposition of classical states or qubits). Our protocol, on the other hand, relies on all four prepared states which are path-spin entangled single particle states. The path-spin entanglement is crucial in security analysis, akin to the E91 protocol, as we show later. Our scheme, based on single particle hybrid entangled states thus relies on a combination elements of both the BB84 and the E91 protocols, but with several important differences.

Our protocol proceeds as follows. Like the BB84 protocol, we also divide the four states into two groups. Each group contains two orthogonal states, but each state of one group is non-orthogonal to the states of the other group. The non-orthogonality is important in the sense that it provides security to the protocol. This follows from an important feature (closely related to no-cloning theorem) of quantum mechanics which states that if a measurement allows one to gain information about the state of a quantum system, then, in general, the state of this quantum system will be disturbed, unless we know in advance that the possible states of the original quantum system are orthogonal to each other. Alice first prepares four path-spin entangled states and places them into two groups. She then randomly selects one group and chooses one of the states from it. Thereafter, she sends that state to Bob. In the next stage, Bob uses two mirrors, phase-shifters and beam splitters. He is free to play with one of the two settings by randomly adjusting the phase-shifter parameter to $\phi = 0$ or $\phi = \frac{\pi}{2}$. Then he performs measurements on the path state vector and the spin state vector using a couple of Stern-Gerlach devices. The measurement on the spin state vector is performed in two different bases. This experiment is repeated many times and each time he notes down the measurement outcome of the path and the spin state vector. In the next step, Alice declares the group (but not the member of the group) she used in each experiment through an authenticated public channel. The existence of an authenticated channel between Alice and Bob, which gives Bob an advantage over Eve, is essential to the security of QKD. After listening to Alice's announcement, Bob informs Alice which data should be kept and which ones should be aborted. The kept data generates the quantum key. This makes the ending of the initial phase.

In the final phase (security checking phase) of our protocol Bob checks whether the generated key is secure. The final phase goes likethe E91 protocol, but with an additional difference. In the E91 protocol, Bob discards some measurement results which are not used in quantum key generation, but in our protocol the aborted measurement outcomes are used in security check. In this way, we can use each and every experiment either in quantum key generation or in security checking. In the security checking, only the aborted cases are considered. In these aborted cases, Alice declares the member of the group, i.e., the prepared intraparticle entangled state through the authenticated public channel. Bob then uses this information to prepare a mixed state and then checks whether the prepared mixed state violates Bell's inequality or not. If the prepared mixed state violates Bell's inequality then the quantum key is considered to be secure, otherwise not. Since this mixed state is a single particle entangled state, the violation of Bell's inequality corresponds in this case to the quantum mechanical violation of non-contextuality. The procedure for demonstrating the violation of contextuality by a single-particle path-spin entangled state was discussed by Basu et al. [15] using a Mach-Zehnder type interferometric set-up, and the actual experiment using single neutrons was performed by Rauch et al. [16]. Thus, the security check for the present protocol could be performed practically using this approach.

GENERATION OF INTRAPARTICLE ENTANGLEMENT

Let us consider a spin-1/2 particle (say, a neutron) that has an initial spin polarized state along the $+\hat{z}-axis$ (denoted by $|\uparrow_z\rangle$). Taking into consideration its path (or position) variables, the joint path-spin state can be written as

$$|\psi_0\rangle_{ps} = |\psi_0\rangle_p \otimes |\uparrow_z\rangle_s \tag{1}$$

where the subscripts p and s refer to the path and the spin variables respectively. The ensemble of neutrons corresponding to $|\psi_0\rangle_{ps}$ are with Alice, incident on a beam-splitter (BS1) whose transmission and reflection probabilities are $|\alpha|^2$ and $|\beta|^2$ respectively. Any given incident particle can then emerge along either the reflected or the transmitted channel corresponding to the state designated by $|\psi_R\rangle$ or $|\psi_T\rangle$ respectively. Here we recall that for any given lossless beam-splitter, arguments using the unitarity condition show that for the particles incident on the beam-splitter, the phase shift between the transmitted and the reflected states of particles is essentially $\frac{\pi}{2}$. For simplifying our treatment, we take the transmission and reflection amplitudes of BS1 and BS2 to be real quantities where $\alpha^2+\beta^2=1$. Note that the beam-splitter acts only on the path-states without affecting the spin-state of the particles, while in our argument a crucial role is played by the mutually orthogonal path states $|\psi_R\rangle$ and $|\psi_T\rangle$ which are eigenstates of the projection operators $P(\psi_R)$ and $P(\psi_T)$ respectively. These projection operators can be regarded as corresponding to the observables that pertain to the determination of which channel a particle is found to be in. For example, the results of such a measurement for the reflected (transmitted) channel with binary alternatives are given by the eigenvalues of $P(\psi_R)(P(\psi_T))$; the eigen- value $+1(0)$ corresponds to a neutron being found (not found) in the channel represented by $|\psi_R\rangle(P(\psi_T))$. The state of a particle emergent from BS1 can then be written as

$$|\psi_0\rangle_{ps} \rightarrow |\psi_1\rangle_{ps} = (\alpha|\psi_T\rangle_p + i\beta|\psi_R\rangle_p) \otimes |\uparrow_z\rangle_s \tag{2}$$

The state vectors $|\psi_T\rangle_p$, $|\psi_R\rangle_p$, $|\uparrow_z\rangle_s$, and $|\downarrow_z\rangle_s$ can be represented as

$$\begin{aligned} &|\psi_T\rangle_p \equiv \begin{pmatrix} 0 \\ 1 \end{pmatrix}, |\psi_R\rangle_p \equiv \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ &|\uparrow_z\rangle_s \equiv |0\rangle_s \equiv \begin{pmatrix} 0 \\ 1 \end{pmatrix}, |\downarrow_z\rangle_s \equiv |1\rangle_s \equiv \begin{pmatrix} 1 \\ 0 \end{pmatrix} \end{aligned} \tag{3}$$

Next, let us suppose that the particles in the channel corresponding to $|\psi_T\rangle_p$ pass through a spin-flipper (SF) (that contains a uniform magnetic field along, say, the $\hat{x}-axis$) which flips the state $|\uparrow_z\rangle_s$ to $|\downarrow_z\rangle_s$. As a consequence of this insertion of a spin-flipper in one of the channels, the particle 1 with Alice has now the path-spin entangled state given by

$$\begin{aligned} |\psi\rangle_{ps} &= \alpha|\uparrow_z\rangle_s \otimes |\psi_T\rangle_p + i\beta|\downarrow_z\rangle_s \otimes |\psi_R\rangle_p \\ &\equiv \alpha|0\rangle_s \otimes |\psi_T\rangle_p + i\beta|1\rangle_s \otimes |\psi_R\rangle_p \end{aligned} \tag{4}$$

QUANTUM KEY DISTRIBUTION SCHEME AND ITS SECURITY

In this section, we discuss our quantum key distribution scheme. Our protocol goes as follows.

A. Intraparticle entanglement prepared by Alice

Let us consider the four intraparticle entangled states $|\Psi\rangle$, $|\Psi_\perp\rangle$, $|\Phi\rangle$, $|\Phi_\perp\rangle$, where

$$\begin{aligned}
|\Psi\rangle &= \frac{1}{\sqrt{2}}(|0\rangle_s \otimes |\psi_T\rangle_p + i|1\rangle_s \otimes |\psi_R\rangle_p)\\
|\Psi_\perp\rangle &= \frac{1}{\sqrt{2}}(|0\rangle_s \otimes |\psi_T\rangle_p - i|1\rangle_s \otimes |\psi_R\rangle_p)\\
|\Phi\rangle &= \frac{1}{\sqrt{2}}(\frac{|0\rangle_s + |1\rangle_s}{\sqrt{2}} \otimes |\psi_T\rangle_p + i\frac{|0\rangle_s - |1\rangle_s}{\sqrt{2}} \otimes |\psi_R\rangle_p)\\
|\Phi_\perp\rangle &= \frac{1}{\sqrt{2}}(\frac{|0\rangle_s + |1\rangle_s}{\sqrt{2}} \otimes |\psi_T\rangle_p - i\frac{|0\rangle_s - |1\rangle_s}{\sqrt{2}} \otimes |\psi_R\rangle_p)
\end{aligned} \tag{5}$$

The former two intraparticle entangled states $|\Psi\rangle$ and $|\Psi_\perp\rangle$ are prepared in a way described in the previous section while the latter two entangled states $|\Phi\rangle$ and $|\Phi_\perp\rangle$ are generated when the Hadamard gate is applied on the spin state of $|\Psi\rangle$ and $|\Psi_\perp\rangle$. We should note that the states $|\Psi\rangle$ and $|\Psi_\perp\rangle$ are not orthogonal to $|\Phi\rangle$ and $|\Phi_\perp\rangle$. Alice prepares any one of the above four intrapaticle entangled states randomly and then sends that state to Bob. This event is repeated many times.



FIG. 1: A spin-1/2 particle falls on the beam-splitter BS1. A spin-flipper is placed along the transmitted channel, and a phase-shifters applied in the reflected path. Bob uses anothet phase-shifter (PS2) and then recombines the reflected and the transmitted channels. Finally, Bob performs path and spin measurements using the Stern-Gerlach devices SG1 and SG2.

B. Path and Spin measurement by Bob

Bob places the mirrors M1 and M2 along the path $|\psi_T\rangle$ and $|\psi_R\rangle$ and in addition to this, a phase-shifter (PS2) is also kept along the path $|\psi_R\rangle$ that changes the phase of the particle by ϕ. Subsequently, Bob uses a 50:50 beam splitter (BS2), to recombine the beams corresponding to $|\psi_T\rangle$ and $|\psi_R\rangle$. The output path states after interacting with beam splitter are given by

$$\begin{aligned}
|\psi'_T\rangle &= \frac{1}{\sqrt{2}}(|\psi_T\rangle + ie^{i\phi}|\psi_R\rangle)\\
|\psi'_R\rangle &= \frac{1}{\sqrt{2}}(i|\psi_T\rangle + e^{i\phi}|\psi_R\rangle)
\end{aligned} \tag{6}$$

If Alice sends the states $|\Psi\rangle$, $|\Psi_\perp\rangle$, $|\Phi\rangle$ or $|\Phi_\perp\rangle$, then after emerging from beam splitter (BS2), the possible resulting states at Bob's site are given by

$$\begin{aligned}
|\Psi'\rangle &= \frac{1}{\sqrt{2}}(i|\psi'_T\rangle \otimes |\chi_1\rangle + |\psi'_R\rangle \otimes |\chi_2\rangle)\\
|\Psi'_\perp\rangle &= \frac{1}{\sqrt{2}}(|\psi'_T\rangle \otimes |\chi_2\rangle - i|\psi'_R\rangle \otimes |\chi_1\rangle)\\
|\Phi'\rangle &= \frac{1}{\sqrt{2}}(i|\psi'_T\rangle \otimes |\chi_3\rangle + |\psi'_R\rangle \otimes |\chi_4\rangle)\\
|\Phi'_\perp\rangle &= \frac{1}{\sqrt{2}}(i|\psi'_T\rangle \otimes |\chi_4\rangle + |\psi'_R\rangle \otimes |\chi_3\rangle)
\end{aligned} \tag{7}$$

where $|\chi_1\rangle = \frac{1}{\sqrt{2}}(|1\rangle_s - ie^{i\phi}|0\rangle_s)$, $|\chi_2\rangle = \frac{1}{\sqrt{2}}(|1\rangle_s + ie^{i\phi}|0\rangle_s)$, $|\chi_3\rangle = \frac{1}{\sqrt{2}}(\frac{1+e^{i\phi}}{\sqrt{2}}|0\rangle_s + \frac{1-e^{i\phi}}{\sqrt{2}}|1\rangle_s)$, $|\chi_4\rangle = \frac{1}{\sqrt{2}}(\frac{1-e^{i\phi}}{\sqrt{2}}|0\rangle_s + \frac{1+e^{i\phi}}{\sqrt{2}}|1\rangle_s)$.

Let us consider that Bob randomly uses two settings of his apparatus by choosing the value of the phase shifter parameter ϕ, say, $\phi = 0$ and $\phi = \frac{\pi}{2}$. If Bob chooses $\phi = 0$ then eq.(7) reduces to

$$\begin{aligned}
|\Psi^{(0)}\rangle &= \frac{1}{\sqrt{2}}(i|\psi'_T\rangle \otimes |\chi_1^{(0)}\rangle + |\psi'_R\rangle \otimes |\chi_2^{(0)}\rangle)\\
|\Psi_\perp^{(0)}\rangle &= \frac{1}{\sqrt{2}}(|\psi'_T\rangle \otimes |\chi_2^{(0)}\rangle - i|\psi'_R\rangle \otimes |\chi_1^{(0)}\rangle)\\
|\Phi^{(0)}\rangle &= \frac{1}{\sqrt{2}}(i|\psi'_T\rangle \otimes |\chi_3^{(0)}\rangle + |\psi'_R\rangle \otimes |\chi_4^{(0)}\rangle)\\
|\Phi_\perp^{(0)}\rangle &= \frac{1}{\sqrt{2}}(i|\psi'_T\rangle \otimes |\chi_4^{(0)}\rangle + |\psi'_R\rangle \otimes |\chi_3^{(0)}\rangle)
\end{aligned} \tag{8}$$

where $|\chi_1^{(0)}\rangle = \frac{1}{\sqrt{2}}(|1\rangle_s - i|0\rangle_s)$, $|\chi_2^{(0)}\rangle = \frac{1}{\sqrt{2}}(|1\rangle_s + i|0\rangle_s)$, $|\chi_3^{(0)}\rangle = |0\rangle_s$, $|\chi_4^{(0)}\rangle = |1\rangle_s$.

If Bob chooses the setting $\phi = \frac{\pi}{2}$ then he further performs a Hadamard gate transformation on the spin qubit and thereafter eq.(7) reduces to

$$\begin{aligned}
|\Psi^{(\frac{\pi}{2})}\rangle &= \frac{1}{\sqrt{2}}(i|\psi'_T\rangle \otimes |\chi_1^{(\frac{\pi}{2})}\rangle + |\psi'_R\rangle \otimes |\chi_2^{(\frac{\pi}{2})}\rangle)\\
|\Psi_\perp^{(\frac{\pi}{2})}\rangle &= \frac{1}{\sqrt{2}}(|\psi'_T\rangle \otimes |\chi_2^{(\frac{\pi}{2})}\rangle - i|\psi'_R\rangle \otimes |\chi_1^{(\frac{\pi}{2})}\rangle)\\
|\Phi^{(\frac{\pi}{2})}\rangle &= \frac{1}{\sqrt{2}}(i|\psi'_T\rangle \otimes |\chi_3^{(\frac{\pi}{2})}\rangle + |\psi'_R\rangle \otimes |\chi_4^{(\frac{\pi}{2})}\rangle)\\
|\Phi_\perp^{(\frac{\pi}{2})}\rangle &= \frac{1}{\sqrt{2}}(i|\psi'_T\rangle \otimes |\chi_4^{(\frac{\pi}{2})}\rangle + |\psi'_R\rangle \otimes |\chi_3^{(\frac{\pi}{2})}\rangle)
\end{aligned} \tag{9}$$

where $|\chi_1^{(\frac{\pi}{2})}\rangle = |0\rangle_s$, $|\chi_2^{(\frac{\pi}{2})}\rangle = |1\rangle_s$, $|\chi_3^{(\frac{\pi}{2})}\rangle = \frac{1}{\sqrt{2}}(|0\rangle_s - i|1\rangle_s)$, $|\chi_4^{(\frac{\pi}{2})}\rangle = \frac{1}{\sqrt{2}}(|0\rangle_s + i|1\rangle_s)$.

Using his Stern-Gerlach devices Bob makes path measurement in the basis $\{|\psi'_T\rangle, |\psi'_R\rangle\}$. The path state vectors are $|\psi'_T\rangle$ and $|\psi'_R\rangle$ with path observable given by

$$\begin{aligned}\hat{\sigma}^p_\phi &= |\psi'_T\rangle\langle\psi'_T| - |\psi'_R\rangle\langle\psi'_R| \\ &= \begin{pmatrix} 0 & -ie^{-i\phi} \\ ie^{i\phi} & 0 \end{pmatrix} \end{aligned} \tag{10}$$

The path observable $\hat{\sigma}^p_\phi$ varies according to the magnitude of the phase shift ϕ. It has two eigenvalues ± 1. The eigenvalue $+1$ and -1 correspond to the measurement of path state vectors $|\psi'_T\rangle$ and $|\psi'_R\rangle$ respectively. The spin measurement is performed either in the basis $\{|0\rangle_s, |1\rangle_s\}$ or in the basis $\{\frac{1}{\sqrt{2}}(|1\rangle_s - i|0\rangle_s), \frac{1}{\sqrt{2}}(|1\rangle_s + i|0\rangle_s)\}$, which are the eigenstates of the observables σ_y and σ_z respectively, with eigenvalues ± 1.

C. Quantum key distribution scheme

Step-I: Alice prepares the intraparticle entangled states and then divides them into two groups G_1^A and G_2^A, where

$$\begin{aligned} G_1^A &= \{|\Psi\rangle, |\Psi_\perp\}\\ G_2^A &= \{|\Phi\rangle, |\Phi_\perp\} \end{aligned} \tag{11}$$

She then randomly picks up one of the states either from G_1^A or from G_2^A and sends that to Bob.

Step-II: Bob randomly changes his settings by choosing the value of the phase shifter either $\phi = 0$ or $\phi = \frac{\pi}{2}$ irrespective of the state sent by Alice. Bob applies a Hadamard gate transformation on the spin component if he chooses $\phi = \frac{\pi}{2}$ setting and does nothing if the $\phi = 0$ setting is used.

Step-III: Bob measures the path state vector in the basis $\{|\psi'_T\rangle, |\psi'_R\rangle\}$ and also perform measurement on spin component either in the basis $\{|0\rangle_s, |1\rangle_s\}$ or in the basis $\{\frac{1}{\sqrt{2}}(|1\rangle_s - i|0\rangle_s), \frac{1}{\sqrt{2}}(|1\rangle_s + i|0\rangle_s)\}$. Bob then noted down the path and spin measurement outcome.

Step-IV: The experiment described in the above three steps is repeated many times. Alice then publicly declares the group (not the member of the group) from which she chooses the state randomly. After listening to Alice's group declaration, Bob informs Alice which experimental data should be kept and which ones should be aborted. The data which are kept are used in generating the secret key, and the data which are aborted are used in the security check, as described in the table below:

State send by Alice	Bob's settings	Path measurement outcome	Spin measurement outcome	Decision
$\|\Psi\rangle$	$\phi = 0$	$\|\psi'_T\rangle$ $(\|\psi'_R\rangle)$	$\frac{1}{\sqrt{2}}(\|1\rangle_s - i\|0\rangle_s)$ $(\frac{1}{\sqrt{2}}(\|1\rangle_s + i\|0\rangle_s))$	KEEP
		$\|\psi'_T\rangle$ $(\|\psi'_R\rangle)$	$\|0\rangle_s$ $(\|1\rangle_s)$	ABORT
	$\phi = \frac{\pi}{2}$	$\|\psi'_T\rangle$ $(\|\psi'_R\rangle)$	$\frac{1}{\sqrt{2}}(\|1\rangle_s - i\|0\rangle_s)$ $(\frac{1}{\sqrt{2}}(\|1\rangle_s + i\|0\rangle_s))$	ABORT
		$\|\psi'_T\rangle$ $(\|\psi'_R\rangle)$	$\|0\rangle_s$ $(\|1\rangle_s)$	KEEP
$\|\Psi_\perp\rangle$	$\phi = 0$	$\|\psi'_T\rangle$ $(\|\psi'_R\rangle)$	$\frac{1}{\sqrt{2}}(\|1\rangle_s + i\|0\rangle_s)$ $(\frac{1}{\sqrt{2}}(\|1\rangle_s - i\|0\rangle_s))$	KEEP
		$\|\psi'_T\rangle$ $(\|\psi'_R\rangle)$	$\|0\rangle_s$ $(\|1\rangle_s)$	ABORT
	$\phi = \frac{\pi}{2}$	$\|\psi'_T\rangle$ $(\|\psi'_R\rangle)$	$\frac{1}{\sqrt{2}}(\|1\rangle_s - i\|0\rangle_s)$ $(\frac{1}{\sqrt{2}}(\|1\rangle_s + i\|0\rangle_s))$	ABORT
		$\|\psi'_T\rangle$ $(\|\psi'_R\rangle)$	$\|1\rangle_s$ $(\|0\rangle_s)$	KEEP
$\|\Phi\rangle$	$\phi = 0$	$\|\psi'_T\rangle$ $(\|\psi'_R\rangle)$	$\frac{1}{\sqrt{2}}(\|1\rangle_s - i\|0\rangle_s)$ $(\frac{1}{\sqrt{2}}(\|1\rangle_s + i\|0\rangle_s))$	ABORT
		$\|\psi'_T\rangle$ $(\|\psi'_R\rangle)$	$\|0\rangle_s$ $(\|1\rangle_s)$	KEEP
	$\phi = \frac{\pi}{2}$	$\|\psi'_T\rangle$ $(\|\psi'_R\rangle)$	$\frac{1}{\sqrt{2}}(\|1\rangle_s - i\|0\rangle_s)$ $(\frac{1}{\sqrt{2}}(\|1\rangle_s + i\|0\rangle_s))$	KEEP
		$\|\psi'_T\rangle$ $(\|\psi'_R\rangle)$	$\|0\rangle_s$ $(\|1\rangle_s)$	ABORT
$\|\Phi_\perp\rangle$	$\phi = 0$	$\|\psi'_T\rangle$ $(\|\psi'_R\rangle)$	$\frac{1}{\sqrt{2}}(\|1\rangle_s - i\|0\rangle_s)$ $\frac{1}{\sqrt{2}}(\|1\rangle_s + i\|0\rangle_s)$	ABORT
		$\|\psi'_T\rangle$ $(\|\psi'_R\rangle)$	$\|1\rangle_s$ $(\|0\rangle_s)$	KEEP
	$\phi = \frac{\pi}{2}$	$\|\psi'_T\rangle$ $(\|\psi'_R\rangle)$	$\frac{1}{\sqrt{2}}(\|1\rangle_s + i\|0\rangle_s)$ $\frac{1}{\sqrt{2}}(\|1\rangle_s - i\|0\rangle_s)$	KEEP
		$\|\psi'_T\rangle$ $(\|\psi'_R\rangle)$	$\|0\rangle_s$ $(\|1\rangle_s)$	ABORT

D. Security check

Let us consider the data which are not used in the key generation. For example, let us suppose that the experiments are performed 1000 times. Further, consider that 500 times they (Alice and Bob) succeded in generating the key and the remaining 500 times they did not. Now, consider those remaining 500 cases for which Bob conveys the word "ABORT" to Alice. In these cases Alice announces the member of the group (i.e. input state that she sent to Bob) through an authenticated channel. If we consider the situation where Alice announces the state $|\Psi\rangle$ 200 times and the states $|\Psi_\perp\rangle$, $|\Phi\rangle$ and $|\Phi_\perp\rangle$ 100 times each, then the probability of occurence of the state $|\Psi\rangle$ is $\frac{2}{5}$ and the probability of occurence of the states $|\Psi_\perp\rangle$,$|\Phi\rangle$ and $|\Phi_\perp\rangle$ are $\frac{1}{3}$ each. With this information, Bob prepares the mixed state

$$\rho = \frac{2}{5}|\Psi\rangle\langle\Psi| + \frac{1}{5}(|\Psi_\perp\rangle\langle\Psi_\perp| + |\Phi\rangle\langle\Phi| + |\Phi_\perp\rangle\langle\Phi_\perp|), \tag{12}$$

If the prepared mixed state violates Bell's inequality, then the generated key considered to be secure, otherwise the key is not secure. In the above example, one can check that the state described by the density operator satisfies Bell's inequality. Hence the key is not secure.

We now generalize the above described situation and consider all the aborted cases where the states $|\Psi\rangle$, $|\Psi_\perp\rangle$, $|\Phi\rangle$ and $|\Psi_\perp\rangle$ occur with probability p_1, p_2, p_3 and p_4 respectively. We should note that in the case of aborted experiments, Bob determines the value of p_1, p_2, p_3 and p_4 only when Alice declares the sent state. With the value of p_1, p_2, p_3 and p_4, Bob prepares the mixed state

$$\begin{aligned}&\rho^B = p_1|\Psi\rangle\langle\Psi| + p_2|\Psi_\perp\rangle\langle\Psi_\perp| + p_3|\Phi\rangle\langle\Phi| + p_4|\Phi_\perp\rangle\langle\Phi_\perp|,\\ &p_1 + p_2 + p_3 + p_4 = 1\end{aligned} \tag{13}$$

For the security of our generated key, we have to check whether the state described by the density operator ρ^B violates Bell's inequality or not. We use Horodecki's [17] criteria to show the Bell's inequality violation, which states that Bell's inequality is violated iff $M(\rho) > 1$,

$$M(\rho) = \lambda + \mu \tag{14}$$

where λ and μ are the two largest eigenvalues of the symmetric matrix $T^\dagger T$, where T denotes the correlation matrix. The correlation matrix corresponding to the state described by the density operator ρ^B is given by

$$T = \begin{pmatrix} p_1 - p_2 & -(p_3 - p_4) & 0 \\ 0 & p_1 - p_2 & p_3 - p_4 \\ p_3 + p_4 & 0 & -(p_1 + p_2) \end{pmatrix} \tag{15}$$

The symmetric matrix $T^\dagger T$ is given by

$$T^\dagger T = \begin{pmatrix} t_{11} & t_{12} & t_{13} \\ t_{12} & t_{22} & t_{23} \\ t_{13} & t_{23} & t_{33} \end{pmatrix} \tag{16}$$

where $t_{11} = (p_1 - p_2)^2 + (p_3 + p_4)^2$, $t_{12} = -(p_1 - p_2)(p_3 - p_4)$, $t_{13} = -(p_1 + p_2)(p_3 + p_4)$, $t_{22} = (p_1 - p_2)^2 + (p_3 - p_4)^2$, $t_{23} = (p_1 - p_2)(p_3 - p_4)$, $t_{33} = (p_1 + p_2)^2 + (p_3 - p_4)^2$.

Now let us consider a particular case where $p_1 = p$, $p_2 = p_3 = p_4 = \frac{1-p}{3}$. In this case, the two largest eigenvalues are given by

$$\lambda = \frac{16p^2 - 8p + 1}{9}, \mu = \frac{4p^2 - 2p + 1}{3} + \frac{2\sqrt{20p^4 + 30p^2 - 44p^3 + 2 - 8p}}{9} \tag{17}$$

Thus the quantity $M(\rho)$ is given by

$$M(\rho) = \lambda + \mu = \frac{28p^2 - 14p + 4}{9} + \frac{2\sqrt{20p^4 + 30p^2 - 44p^3 + 2 - 8p}}{9} \tag{18}$$

It can be easily shown that the violation occurs only for $p \geq 0.67$. Therefore, we can conclude that if $p \geq 0.67$ then the quantum key distribution scheme is successful, otherwise not.

It might be interesting to estimate the percentage of the data used in the key generation process corresponding to any given value of p for the mixed state. Consider the four states be divided into two basis sets B_1 and B_2, where $B_1 = \{|\Psi\rangle, |\Psi_\perp\}$, and $B_2 = \{|\Phi\rangle, |\Phi_\perp\rangle\}$. Our protocol fails to generate a secure key in the cases: (i) if Alice chooses one of the states from B_1 to send it to Bob, and if Bob uses the phase shifter parameter $\phi = 0$; or (ii) if Alice chooses one of the states from B_2 to send it to Bob and if Bob uses the phase shifter parameter $\phi = \frac{\pi}{2}$. Let η_1 and η_2 be the probabilities of failure for first and second case respectively.

$$\begin{aligned} \eta_1 &= P(\phi = 0/|\Psi\rangle).P(|\Psi\rangle) + P(\phi = 0/|\Psi_\perp\rangle).P(|\Psi_\perp\rangle) \\ &= \frac{1}{2}.p_1 + \frac{1}{2}.p_2 \end{aligned} \tag{19}$$

$$\begin{aligned} \eta_2 &= P(\phi = \frac{\pi}{2}/|\Phi\rangle).P(|\Phi\rangle) + P(\phi = \frac{\pi}{2}/|\Phi_\perp\rangle).P(|\Phi_\perp\rangle) \\ &= \frac{1}{2}.p_3 + \frac{1}{2}.p_4 \end{aligned} \tag{20}$$

Now choosing $p_1 = p$ and $p_2 = p_3 = p_4 = \frac{1-p}{3}$, we have

$$\eta_1 = \frac{2p+1}{6} \tag{21}$$

$$\eta_2 = \frac{1-p}{3} \tag{22}$$

For different values of $p \geq 0.67$, the values of η_1 and η_2 are listed in the table given below:

Sl. No.	p	$M(\rho)$	η_1	η_2
1.	0.67	1.01	0.39	0.11
2.	0.70	1.08	0.40	0.10
3.	0.75	1.20	0.41	0.08
4.	0.80	1.34	0.43	0.06
5.	0.85	1.49	0.45	0.05
6.	0.90	1.64	0.46	0.03
7.	0.95	1.81	0.48	0.01
8.	1.0	2.0	0.50	0.00

CONCLUSIONS

In this work we have formulated a quantum key distribution scheme using intraparticle entangled states. Our protocol proceeds in phases like the BB84 and the E91 protocols, but with notable differences. Our protocol is based on single particle states with two entangled degrees of freedom. The entanglement is crucial for security analysis, which is based on the quantum violation of noncontextuality for single particle mixed entangled states. Further, no data is wasted since the data which is not used in quantum key generation can be used in security checking. Thus, we can use each and every experiment either in a quantum key generation or in security checking. Our protocol provides an example of a practical use of quantum contextuality. Moreover, it shoulc be possible to experimentally realize our scheme, with exisitng set-ups such as the one on neutron interferometry [16]. We conclude by noting that though the scheme presented here is based on the path-spin entanglement of spin-1/2 particles, it can be easily extended for other single particle entangled states, such as using the path and polarization variables of photons, for instance.

Acknowledgements: ASM and DH acknowledge support from the DST Project SR/S2/PU-16/2007. DH also thanks the Centre for Science, Kolkata for support.

ACKNOWLEDGEMENT

---

[*] tapisatya@gmail.com
[†] dhome@bosemain.boseinst.ac.in
[‡] archan@bose.res.in
[§] apan@bosemain.boseinst.ac.in

[1] N. Gisin, G. Ribordy, W. Tittel, and H. Zbinden, Rev. Mod. Phys. **74**, 145 (2002).

[2] W. K. Wootters, and W. H. Zurek, Nature (London) **299**, 802 (1982).

[3] C. H. Bennett, and G. Brassard, in Proceedings of the IEEE International Conference on Computers, Systems and Signal Processing, Bangalore, India, (IEEE, New York), 175 (1984).

[4] M. Lucamarini, and S. Mancini, Phys. Rev. Lett. **94**, 140501 (2005).

[5] D. Bruss, Phys. Rev. Lett. 81, 3018 (1998).

[6] X. B. Wang, Phys. Rev. Lett. **94**, 230503 (2005).

[7] B. Kraus, N. Gisin, and R. Renner, Phys. Rev. Lett. **95**, 080501 (2005).

[8] H. -K. Lo, H. F. Chau, and M. Ardehali, J. Cryptology **18**, 133 (2005).

[9] A. Acin, N. Gisin, and L. Masanes, Phys. Rev. Lett. 97, 120405 (2006).

[10] Y. Adachi, T. Yamamoto, M. Koashi, and N. Imoto, Phys. Rev. Lett. 99, 180503 (2007).

[11] A. K. Ekert, Phys. Rev. Lett. **67**, 661 (1991).

[12] J. S. Bell, Physics 1 (Long Island City, N.Y.), 195 (1964).

[13] T. Jennewein, C. Simon, G. Weihs, H. Weinfurter, and A. Zeilinger, Phys. Rev. Lett. 84, 4729 (2000).

[14] A. Ling, M. P. Peloso, I. Marcikic, V. Scarini, A. Lamaslinares, and C. Kurtsiefer, Phys. Rev. A 78, 020301 (R) (2008).

[15] S. Basu, S. Bandyopadhyay, G. Kar, and D. Home, Phys. lett. A 279, 281 (2001).

[16] Y. Hasegawa, R. Loidl, G. Badurek, M. Baron, and H. Rauch, Nature 425, 45 (2003).

[17] R. Horodecki, M. Horodecki, and P. Horodecki, Phys. Lett. A 222, 21 (1996).